

## Seminar 2

### (S2.1)

- (i)  $(X, 1_X)$  is minimal if and only if  $|X| = 1$ .
- (ii) If  $(X, T)$  is minimal, then  $T$  is surjective.
- (iii) A factor of a minimal TDS is also minimal.
- (iv) If a product TDS is minimal, then so are each of its components.
- (v) If  $(X_1, T_{X_1})$ ,  $(X_2, T_{X_2})$  are two minimal subsystems of a TDS  $(X, T)$ , then either  $X_1 \cap X_2 = \emptyset$  or  $X_1 = X_2$ .
- (vi) A disjoint union of two TDSs is never a minimal TDS.

*Proof.* (i) Remark that for all  $x \in X$ ,  $\overline{\text{orb}_+(x)} = \{x\}$ .

- (ii) By Corollary 1.3.3.5, there exists a nonempty closed set  $B \subseteq X$  such that  $T(B) = B$ . Since  $(X, T)$  is minimal, we must have  $B = X$ .
- (iii) Let  $(X, T)$  be minimal and  $\varphi : (X, T) \rightarrow (Y, S)$  be a surjective homomorphism. Assume  $\emptyset \neq A \subseteq Y$  is a nonempty closed  $S$ -invariant subset of  $Y$ . We have to prove that  $A = Y$ . Let  $B := \varphi^{-1}(A) \subseteq X$ . Then  $B$  is closed and nonempty, since  $\varphi$  is continuous and surjective. Furthermore,

$$\begin{aligned} T(B) &= T(\varphi^{-1}(A)) = \{Tx \mid \varphi(x) \in A\} \subseteq \{Tx \mid (S \circ \varphi)(x) \in A\} \\ &\quad \text{since } (S \circ \varphi)(x) \in S(A) \subseteq A \\ &= \{Tx \mid (\varphi \circ T)(x) \in A\} \subseteq \varphi^{-1}(A) = B. \end{aligned}$$

Thus,  $B$  is a nonempty closed  $T$ -invariant subset of  $X$ , so we must have  $B = X$ . Using again the surjectivity of  $\varphi$ , it follows that

$$Y = \varphi(X) = \varphi(\varphi^{-1}(A)) = A.$$

(iv) By (iii) and Proposition 1.3.4.1.(ii).

(v) We have that  $X_1, X_2$  are nonempty closed  $T$ -invariant subsets of  $X$ . Let  $Y := X_1 \cap X_2$ . Then  $Y$  is a closed  $T_{X_1}$ -invariant subset of  $X_1$  (resp. a closed  $T_{X_2}$ -invariant subset of  $X_2$ ), hence from minimality we must have  $Y = \emptyset$  or  $Y = X_1 = X_2$ .

(vi) By Lemma 1.3.5.2.(i). □

**(S2.2)** Let  $(X, T)$  be a TDS and assume that  $X$  is metrizable. For any  $x \in X$ , the following are equivalent:

(i)  $x$  is recurrent.

(ii)  $\lim_{k \rightarrow \infty} T^{n_k} x = x$  for some sequence  $(n_k)$  in  $\mathbb{Z}_+$ .

(iii)  $\lim_{k \rightarrow \infty} T^{n_k} x = x$  for some sequence  $(n_k)$  in  $\mathbb{Z}_+$  such that  $\lim_{k \rightarrow \infty} n_k = \infty$ .

*Proof.* (iii)  $\Rightarrow$  (ii) Obviously.

(ii)  $\Rightarrow$  (i) Let  $U$  be an open neighborhood of  $x$ . Since  $\lim_{k \rightarrow \infty} T^{n_k} x = x$ , there exists  $K \in \mathbb{Z}_+$  such that  $T^{n_k} x \in U$  for all  $k \geq K$ .

(i)  $\Rightarrow$  (iii) Use the fact that  $x$  is infinitely recurrent, by Proposition 1.6.0.16. Then  $S_k := rt(x, B_{1/k}(x))$  is an infinite set for every  $k \geq 1$ . Define  $n_1 := \min S_1$ ,  $n_{k+1} := \min S_{k+1} \setminus \{n_k\}$ . Then  $(n_k)$  is a strictly increasing sequence of positive integers, so  $\lim_{k \rightarrow \infty} n_k = \infty$ . Furthermore,  $d(x, T^{n_k} x) < 1/k$  for all  $k \geq 1$ , hence  $\lim_{k \rightarrow \infty} T^{n_k} x = x$ . □

**(S2.3)**

(i) If  $\varphi : (X, T) \rightarrow (Y, S)$  is a homomorphism of TDSs and  $x \in X$  is recurrent (almost periodic) in  $(X, T)$ , then  $\varphi(x)$  is recurrent (almost periodic) in  $(Y, S)$ .

(ii) If  $(A, T_A)$  is a subsystem of  $(X, T)$  and  $x \in A$ , then  $x$  is recurrent (almost periodic) in  $(X, T)$  if and only if  $x$  is recurrent (almost periodic) in  $(A, T_A)$ .

*Proof.* (i) Let  $V$  be an open neighborhood of  $\varphi(x)$ . Since  $\varphi$  is continuous, there exists an open neighborhood  $U$  of  $x$  such that  $\varphi(U) \subseteq V$ .

(a) As  $x$  is recurrent in  $(X, T)$ , we have that  $T^n x \in U$  for some  $n \geq 1$ . We get that

$$S^n(\varphi(x)) = \varphi(T^n x) \in \varphi(U) \subseteq V.$$

It follows that  $\varphi(x)$  is recurrent in  $(X, T)$ .

- (b) As  $x$  is almost periodic in  $(X, T)$ , we have that there exists  $N \geq 1$  such that for all  $m \geq 1$  there exists  $k \in [m, m + N]$  such that  $T^k x \in U$ . We get that

$$S^k(\varphi(x)) = \varphi(T^k x) \in \varphi(U) \subseteq V.$$

It follows that  $\varphi(x)$  is almost periodic in  $(X, T)$ .

- (ii)  $\Leftarrow$  Use (i) and the fact the inclusion  $j_A : (A, T_A) \rightarrow (X, T)$  is a homomorphism.  
 $\Rightarrow$  If  $U$  is an open neighborhood of  $x$  in  $A$ , then  $U = A \cap V$ , where  $V$  is an open neighborhood of  $x$  in  $X$ .

- (a) If  $x$  is recurrent in  $(X, T)$ , we have that  $T^n x \in V$  for some  $n \geq 1$ . It follows that  $T_A^n x = T^n x \in A \cap V = U$ . Thus,  $x$  is recurrent in  $(A, T_A)$ .  
(b) If  $x$  is almost periodic in  $(X, T)$ , we have that there exists  $N \geq 1$  such that for all  $m \geq 1$  there exists  $k \in [m, m + N]$  such that  $T^k x \in V$ . Conclude as above that  $T_A^k x = T^k x \in U$ . Thus,  $x$  is almost periodic in  $(A, T_A)$ .

□

**(S2.4)** Let  $(X, T)$  be a TDS and  $x \in X$ . The following are equivalent:

- (i)  $x$  is almost periodic.  
(ii) For any open neighborhood  $U$  of  $x$ , there exists  $N \geq 1$  such that

$$\text{orb}_+(x) \subseteq \bigcup_{k=0}^N T^{-k}(U).$$

- (iii)  $(\overline{\text{orb}_+(x)}, T_{\overline{\text{orb}_+(x)}})$  is a minimal subsystem.

*Proof.* (i)  $\Rightarrow$  (ii) We have obviously that  $x \in U = T^0(U)$ , so let  $T^m x$  with  $m \geq 1$ . Since  $rt(x, U)$  is syndetic, it follows that there exists  $N \geq 1$  such that  $rt(x, U) \cap [m, m + M] \neq \emptyset$  for all  $m \geq 1$ . Thus, there exists  $p \in [m, m + N]$  such that  $T^p x \in U$ . Letting  $k := p - m \in [0, N]$ , we get that  $T^k(T^m x) = T^p x \in U$ , hence  $T^m x \in T^{-k}(U)$ .

(ii)  $\Rightarrow$  (iii) We shall prove that  $\text{orb}_+(y)$  is dense in  $\overline{\text{orb}_+(x)}$  for every  $y \in \overline{\text{orb}_+(x)}$ , and then apply Proposition 1.5.0.11 to conclude minimality. It suffices to show that  $x \in \overline{\text{orb}_+(y)}$ . Let  $U$  be an open neighborhood of  $x$ . Then, by B.9.0.29.(i), there exists an open neighborhood  $V$  of  $x$  such that  $\overline{V} \subseteq U$ . By (ii), we have an  $N \geq 1$  such that

$$\text{orb}_+(x) \subseteq \bigcup_{k=0}^N T^{-k}(V) \subseteq \bigcup_{k=0}^N T^{-k}(\overline{V}).$$

It follows that

$$y \in \overline{\text{orb}_+(x)} \subseteq \bigcup_{k=0}^N T^{-k}(\overline{V}) \subseteq \bigcup_{k=0}^N T^{-k}(U).$$

This implies  $T^k y \in U$  for some  $k = 0, \dots, N$ . Thus,  $\text{orb}_+(y) \cap U \neq \emptyset$  for any open neighborhood  $U$  of  $x$ , that is  $x \in \overline{\text{orb}_+(y)}$ .

- (iii)  $\Rightarrow$  (i) Apply Proposition 1.6.0.22.

□

**Definition .** A TDS  $(X, T)$  is said to be **isometric** if there exists a metric  $d$  on  $X$  inducing the topology of  $X$  such that  $T$  is an isometry with respect to  $d$ .

(S2.5) Give examples of isometric TDSs.

*Proof.* (i) The rotation on the unit circle  $(\mathbb{S}^1, R_a)$  (see Example 1.1.4 in the lecture).

(ii) Any finite invertible TDS  $(X, T)$ , where  $X$  is a finite metric space with the discrete metric and  $T$  is a bijective mapping. □

(S2.6) Let  $(X, T)$  be an isometric TDS. Then

(i)  $(X, T)$  is minimal if and only if it is forward transitive.

(ii) For every  $x \in X$ ,  $(\overline{\text{orb}_+(x)}, T_{\overline{\text{orb}_+(x)}})$  is a minimal subsystem. Conclude that every point  $x \in X$  is contained in a unique minimal subsystem and that  $(X, T)$  is a disjoint union of minimal subsystems.

(iii) Every point  $x \in X$  is almost periodic.

*Proof.* (i) Assume that  $(X, T)$  is forward transitive, and let  $x_0$  be a forward transitive point. We shall prove that every point  $y \in X$  is forward transitive. It suffices to show that  $x_0 \in \overline{\text{orb}_+(y)}$ .

For every  $\varepsilon > 0$ ,  $\text{orb}_+(x_0) \cap B_\varepsilon(y) \neq \emptyset$ , hence there exists  $m \geq 0$  such that  $d(T^m x_0, y) < \varepsilon/2$ . Since  $X$  is compact, the sequence  $(T^{mn} x_0)_{n \geq 1}$  has a convergent subsequence  $(T^{mn_k} x_0)_{k \geq 1}$ . As  $T^p$  is an isometry for all  $p \geq 0$ , we get that

$$d(x_0, T^{m(n_{k+1}-n_k)} x_0) = d(T^{mn_k} x_0, T^{mn_{k+1}} x_0) \rightarrow 0 \text{ for } k \rightarrow \infty.$$

Hence, there exists  $K \geq 1$  such that  $d(x_0, T^{m(n_{K+1}-n_K)} x_0) < \varepsilon/2$ . Let  $p := m(n_{K+1} - n_K) \geq m$ . It follows that

$$\begin{aligned} d(T^{p-m} y, x_0) &\leq d(T^{p-m} y, T^p x_0) + d(T^p x_0, x_0) = d(y, T^m x_0) + d(T^p x_0, x_0) \\ &< \varepsilon/2 + \varepsilon/2 = \varepsilon. \end{aligned}$$

Thus,  $\text{orb}_+(y) \cap B_\varepsilon(x_0) \neq \emptyset$  for all  $\varepsilon > 0$ . That is,  $x_0 \in \overline{\text{orb}_+(y)}$ .

(ii) For every  $x \in X$ ,  $(\overline{\text{orb}_+(x)}, T_{\overline{\text{orb}_+(x)}})$  is an isometric subsystem of  $(X, T)$  which is also forward transitive, by Lemma 1.4.0.5. Apply (i) to conclude that it is a minimal subsystem containing  $x$ . Uniqueness of the decomposition follows from Proposition 1.5.0.10.(v).

(iii) By (ii) and Proposition 1.6.0.22. □